

## University of Groningen

### **The moonlighting function of pyruvate carboxylase resides in the non-catalytic end of the TIM barrel**

Huberts, Daphne H. E. W.; Venselaar, Hanka; Vriend, Gert; Veenhuis, Marten; van der Klei, Ida J.

*Published in:*  
Biochimica et Biophysica Acta-Molecular Cell Research

*DOI:*  
[10.1016/j.bbamcr.2010.03.018](https://doi.org/10.1016/j.bbamcr.2010.03.018)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2010

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Huberts, D. H. E. W., Venselaar, H., Vriend, G., Veenhuis, M., & van der Klei, I. J. (2010). The moonlighting function of pyruvate carboxylase resides in the non-catalytic end of the TIM barrel. *Biochimica et Biophysica Acta-Molecular Cell Research*, 1803(9), 1038-1042.  
<https://doi.org/10.1016/j.bbamcr.2010.03.018>

#### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

#### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*




Fig. S1. Alignment of pyruvate carboxylases from the methylotrophic yeast *Hansenula polymorpha* (HpPyc1), *Pichia pastoris* (PpPyc1), *Pichia methanolica* (PmPyc1), and *Candida boidinii* (CbPyc1) and both pyruvate carboxylases from *Saccharomyces cerevisiae* (ScPyc1 and ScPyc2). The aligned region contains a large part of the TC domain and the linker region, which links the BC to the TC domain. The three residues (arrows) that are important for the moonlighting function of Pyc1 are conserved in the methylotrophic species but not in *S. cerevisiae*.